

Nausea and Vomiting

In the event of nausea and/or vomiting following surgery, do not take anything by mouth for at least an hour including the prescribed medicine. You should then sip on coke, tea or ginger ale. You should sip slowly over a fifteen-minute period. When the nausea subsides you can begin taking solid foods and the prescribed medicine. If you have persistent nausea and/or vomiting, please contact our office and an anti-nausea medication can be prescribed in most cases.

Other Complications

Expect numbness of the lip, chin, and/or tongue for approximately 5 hours following surgery. This is the effect of the local anaesthetic used during surgery. If it persists, there is no cause for alarm. As stated before surgery, this is usually temporary in nature. You should be aware that if your lip or tongue is numb, you could bite it and not feel the sensation. So be careful and avoid excessively hot or cold food or drinks. Call Dr Wong if you have any questions.

Slight elevation of temperature immediately following surgery is not uncommon. If the fever persists, notify the office. The Panadol and/or Ibuprofen taken for pain will also help to reduce the fever.

Occasionally, patients may feel hard projections in the mouth with their tongue. They are not roots, but they are the bony walls which supported the tooth. These projections usually smooth out spontaneously. If not, they can be removed by the doctor.

If the corners of your mouth are stretched, they may dry out and crack. Your lips should be kept moist with an ointment such as Vaseline.

Sore throat and pain when swallowing are not uncommon. The muscles sometimes swell after surgery. The normal act of swallowing can then become painful. This will subside in 2-3 days.

Finally

Sutures may be placed in the area of surgery to minimize post-operative bleeding and to help healing. Sometimes they become dislodged – this is no cause for alarm. Just remove the suture from your mouth and discard it. The remaining sutures will dissolve spontaneously after about 2-3 weeks.

There will be a hole where the tooth was removed. The hole will gradually close over in the next month and fill in with the new tissue. Keep this area clean as described earlier.

Your case is individual, no two mouths are alike. Do not accept well intended advice from friends. Discuss your problem with the persons best able to effectively help you: the surgeon or your family dentist.

Post operative visit

A post operative visit to see me will be arranged for you. If non-dissolving sutures have been used these will be removed at this visit. Work certificates can also be provided upon request or by phoning the rooms.

If you have any problems or concerns, please call.

SPECIFIC INSTRUCTIONS:

Tooth Exposures

If you have had a tooth exposed there may be a small pack sutured over the tooth that will fall out several days after surgery.

Dental Implants

Implants are often inserted in two stages. The first stage involves insertion of the implant into the bone. If required, a second stage procedure is performed several months later to uncover the implant and place a healing abutment which protrudes through the gum.


Sinus Surgery

If you have had surgery close to or involving the sinus cavity, you will need to limit nose blowing, smoking, sucking on straws and blowing of wind instruments for 2 weeks.

Dr Jaclyn Wong

MBBS BSc (Hons) Post Grad Dip Surg Anat

Minor Oral Surgery



INSTRUCTIONS FOLLOWING SURGERY

Name: _____

Post operative appointment:

- 6 Bond St, Sth Yarra
- 1007 Malvern Rd, Toorak
- 516 Centre Rd, Bentleigh
- 801 Toorak Rd, Hawthorn East
- 1 Thurmond Crt, Endeavour Hills

ALL ENQUIRIES TO:

info@wisdomteethandimplants.com.au

**T: 9882 0839
0411 864 255
0425 755 500**

After Tooth Extraction

The removal of impacted teeth is a surgical procedure. Post-operative care is very important. Unnecessary pain and the complications of infection and swelling can be minimized if these instructions are followed carefully.

Immediately Following Surgery:

- The gauze pad placed over the surgical area should be kept in place for one hour. Upon discharge from hospital, the gauze pad will be removed and you will be provided with spare gauze in case of ongoing bleeding.
- Vigorous mouth rinsing, sucking, spitting or touching the wound area following surgery should be avoided. This may initiate bleeding by causing the blood clot that has formed to become dislodged.
- Take the prescribed pain medications as soon as possible, ideally before the local anaesthetic has worn off.
- Restrict your activities the day of surgery and resume normal activity when you feel comfortable.
- Place ice packs to the sides of your face where surgery was performed. Refer to the section on swelling for explanation.

Bleeding

Bleeding is normally fully controlled intra-operatively, however a certain amount of bleeding may still occur following surgery. Slight bleeding, oozing, or redness in the saliva is common. Excessive bleeding may be controlled by first rinsing or wiping any old clots from your mouth. Then place a gauze pad over the area and bite firmly for thirty minutes. Repeat if necessary. If bleeding continues, bite on a moistened tea bag for thirty minutes. The tannic acid in the tea bag helps to form a clot by contracting bleeding vessels. To minimize further bleeding, do not become excited, sit upright, and avoid exercise. If bleeding does not subside, call immediately for further instructions.

Swelling and Bruising

The amount of expected swelling differs from one procedure to the next. Swelling around the mouth, cheeks, eyes and sides of the face is not uncommon. This is the body's normal reaction to surgery and eventual repair. The swelling will not become apparent until the day following surgery, and will not reach its maximum until 2-3 days post-operatively. However, the swelling may be minimized by the immediate use of ice packs. Two bags filled with ice, or ice packs should be applied to the sides of the face where surgery was performed. The ice packs should be left on for 20 minutes and then removed for 20 minutes while you are awake. After 48 hours, ice has no beneficial effect. If swelling or jaw stiffness has persisted for several days, there is no cause for alarm. This is a normal reaction to surgery. 48 hours following surgery, the application of moist heat to the sides of the face is beneficial in reducing the size of the swelling. Gentle opening and closing of the mouth can help to restore normal movement. After 2-3 days there may be some discolouration of the skin, first blue-black in colour, then turning yellow and disappearing in a few days. This is due to blood spreading beneath the tissues and is a normal post-operative occurrence. Moist heat applied to the area may speed up the removal of the discoloration.

Pain

Pain is a normal consequence of surgery and is expected. Take the medication prescribed as directed and avoid aspirin unless it has been medically prescribed and you have discussed it with Dr Wong. One or two tablets of Panadeine Forte may be taken every four- six hours (maximum 8 tablets per day) in combination with two 200 mg tablets or one 400mg tablet of Ibuprofen every 8 hours. For the first 1-2 days after surgery, it is often effective to alternate the Panadeine Forte and Ibuprofen dosing every 2 hours during waking hours.

The prescribed pain medicine may make you groggy and slow down your reflexes. Do not drive an automobile or work around machinery. Avoid alcoholic beverages. Pain or discomfort following surgery should subside more each day. If pain persists, it may require attention and you should call the office.

Diet

After general anaesthetic or I.V. sedation, room temperature liquids should be initially taken. Do not use straws. Drink from a glass. The sucking motion can cause more bleeding by dislodging the blood clot. You may eat anything soft by chewing away from the surgical sites. Avoid alcohol and smoking for at least 24-48 hours. High calorie, high protein intake is very important. Nourishment should be taken regularly. You should prevent dehydration by taking fluids regularly. Your food intake will be limited for the first few days. You should compensate for this by increasing your fluid intake. At least 5-6 glasses of liquid should be taken daily. Try not to miss a single meal. You will feel better, have more strength, less discomfort and heal faster if you continue to eat. Caution: If you suddenly sit up or stand from a lying position you may become dizzy. If you are lying down following surgery, make sure you sit for one minute before standing.

Keep the Mouth Clean

No rinsing of any kind should be performed until the day following surgery. You can brush your teeth and unaffected areas the night of surgery but rinse gently. The day after surgery, you should rinse your mouth gently after meals with the prescribed mouth rinse (0.2% chlorhexidine). Five days after surgery you must begin irrigating your lower extraction sites (if open) after eating, with the syringe and the mouth rinse or a cup of warm water mixed with a teaspoon of salt. This is essential to keep food and debris from delaying the healing of these areas or causing post-operative infections. Ideally, use the 0.2% chlorhexidine during the first week then change to salt water and continue this until healing of the sockets has occurred.

Antibiotics

If you have been placed on antibiotics, take the tablets or liquid as directed. Antibiotics to prevent infection are only prescribed in patients with a history of recurrent infection, immunosuppression or other specific indications. They are not routinely prescribed in healthy patients unless there is active infection. Discontinue antibiotic use in the event of a rash or other unfavourable reaction. Call the office if this occurs or you have any questions.